



GIG
CYMRU
NHS
WALES

Bwrdd Iechyd Prifysgol
Bae Abertawe
Swansea Bay University
Health Board

Cadeirydd/Chair: **Emma Woollett**
Prif Weithredwr/Chief Executive: **Mark Hackett**

gofalu am ein gilydd, cydweithio, gwella bob amser
caring for each other, working together, always improving

Rydym yn croesawu gohebiaeth yn y Gymraeg neu'r Saesneg. Atebir gohebiaeth Gymraeg yn y Gymraeg, ac ni fydd hyn yn arwain at oedi.
We welcome correspondence in Welsh or English. Welsh language correspondence will be replied to in Welsh, and this will not lead to a delay.

Cais Rhyddid Gwybodaeth / Freedom of Information request **Ein Cyf / Our Ref: 22-I-025**

You asked:

Please complete the questionnaire regarding access to and use of MRI in the diagnosis of axial Spondylarthritis.

Please see below.



Pencadlys BIP Bae Abertawe, Un Porthfa Talbot, Port Talbot, SA12 7BR / Swansea Bay UHB Headquarters, One Talbot Gateway, Port Talbot, SA12 7BR

Bwrdd Iechyd Prifysgol Bae Abertawe yw enw gweithredu Bwrdd Iechyd Lleol Prifysgol Bae Abertawe
Swansea Bay University Health Board is the operational name of Swansea Bay University Local Health Board

UK Radiologist MRI axial SpA FOI
***FOI to assess access to; use of MRI and protocols in the diagnosis
of axial Spondylarthritis***

1. Which Health Board?

Swansea Bay

Part 1 – Access to MRI

2. How many MRI scanners suitable for scanning the spine and SIJ do you have access to?

☐ None

☐ 1

☐ 2

☒ 3

☐ 4 or more

3. What type of scanner do you have access to?

☒ 1.5T

☐ 3T

☐ Other (please state)

4. Roughly how long would an outpatient wait for an MRI in your Health board?

☐ Less than 2 weeks

☐ 2-4 weeks

☒ 1-2 months

☐ 2-3 months

☐ Greater than 3 months

5. Do you have regular meetings or discussions with your rheumatology colleagues?

☒ Weekly

☐ Fortnightly

- ☐ Monthly
- ☐ Quarterly
- ☐ As required
- ☐ Never

6. How is axial SpA MRI imaging reported in your Health Board?

- ☒ Internally by a specialist MSK radiologists
- ☐ Internally by a non-specialist radiologist
- ☒ Outsourced to a specialist MSK radiologist service
- ☐ Outsourced to a non-specialist radiologist service
- ☐ Other (please state)

Part 2 – Use of MRI in diagnosis of axial SpA

7. Are you familiar with the rheumatological term axial Spondyloarthritis (axial SpA)?

- ☒ Yes
- ☐ No

8. Are you familiar with the BRITSpA consensus guidance on MRI for the diagnosis of axial Spondyloarthritis? ([BRITSpA guidance](#))

- ☐ Yes
- ☒ No

9. In what circumstances would you use MRI in the assessment/diagnosis of spondyloarthritis?

- ☐ We do not, or only rarely, use MRI for assessment/diagnosis of spondyloarthritis
- ☒ We only use MRI if the x-rays of the SIJ and spine are normal/not diagnostic
- ☐ We use MRI as a diagnostic test, irrespective of whether there is an abnormality on plain x-rays
- ☐ Other reasons.....

Part 3 – MRI protocols

10. Please write in text below what your standard MRI protocol for the assessment of spondyloarthritis is (specifying (i) field strength, (ii) sequences, (iii) anatomical coverage and (iii) acquisition planes for each element of the protocol):

Sagittal T1 and STIR of the whole spine
Axial T2 of any degenerate levels
Sagittal T1 and PDFS, Axial PDFS of both SI Joints

11. What is the approximate scan time for this protocol? If multiple scanners are used, please give an average or range.

50 minutes

12. When assessing patients for possible early spondyloarthritis with MRI would you routinely scan?

- ☐ Sacroiliac joints only
- ☐ Sacroiliac joints and lumbar spine
- ☐ Sacroiliac joints and thoracic spine
- ☐ Sacroiliac joints and thoracolumbar spine
- ☒ Sacroiliac joints and whole spine
- ☐ Sacroiliac joints and any other spinal segment, including whole spine, as requested by the rheumatologist
- ☐ Other.....

13. If you do not scan any of the spine in the assessment of spondyloarthritis, why not?

- ☐ Spinal features are not necessary in the diagnosis of axial-SpA
- ☐ There are no specific lesions for spondyloarthritis in the spine

- ☐ The scan time is too long
- ☐ It costs too much money to scan
- ☐ Other.....

14.If you do not MRI scan the whole spine in the assessment of spondyloarthritis, why not?

- ☐ Imaging the lumbar / thoracolumbar spine is sufficient to assess spinal features of axial-SpA
- ☐ Spinal features are not necessary in the diagnosis of axial-SpA
- ☐ It takes too long to scan the spine
- ☐ It costs too much money to scan the whole spine
- ☐ Other.....

15.If you are MRI scanning part of, or the whole spine in the assessment of spondyloarthritis would you perform?

- ☒ Sagittal, axial and coronal plane acquisitions
- ☐ Sagittal and axial plane acquisitions only
- ☐ Sagittal plane acquisitions only
- ☐ Other.....

16.If you are MRI scanning the sacroiliac joints, would you perform?

- ☐ Semi-coronal plane acquisitions only
- ☒ Semi-coronal and semi-axial acquisitions

17.When MRI scanning for the assessment of spondyloarthritis which sequences do you use in your protocol? Please tick all that apply.

- ☒ Fat-suppressed, water sensitive sequence (e.g. STIR, T2w imaging with fat suppression or T2w Dixon imaging)
- ☒ Fat-sensitive sequence (T1w imaging)
- ☐ Gadolinium-enhanced imaging
- ☐ Conventional T2w imaging (without fat suppression)

- ☐ Gradient echo imaging (including VIBE or Dixon)
- ☐ Other.....

18. When MRI scanning for the assessment of spondyloarthritis do you perform gadolinium-enhanced imaging of the sacroiliac joints?

- ☐ Yes
- ☒ No

19. When MRI scanning for the assessment of spondyloarthritis do you perform gadolinium-enhanced imaging of the spine?

- ☐ Yes
- ☒ No

Part 4 – MRI lesions and definitions

20. Are you aware of formal recommendations regarding which imaging features should contribute to the identification of a positive MRI of the sacroiliac joints in spondyloarthritis?

- ☒ Yes, please detail below:
- ☒ No

21. What MRI SIJ spondyloarthritis features do you use to make a diagnosis of Spondyloarthritis?

- ☐ Subchondral Bone marrow oedema / osteitis only
- ☐ Erosions
- ☐ Fat infiltration
- ☐ Sclerosis
- ☐ Joint space widening / effusion
- ☐ Enthesitis
- ☐ Capsulitis

- ☐ Fat deposition in the joint space ("backfill")
- ☐ Synovitis
- ☐ New bone formation (areas of ankylosis)
- ☐ All of the above
- ☒ Combination of the pathologies ticked above
- ☐ Other.....

22. Are you aware of formal recommendations regarding which imaging features should contribute to the identification of positive MRI of the spine in spondyloarthritis?

- ☐ Yes, please detail below:
- ☒ No

23. What MRI spinal spondyloarthritis features do you use to make a diagnosis of Spondyloarthritis?

- ☒ Vertebral corner bone marrow oedema
- ☐ Endplate oedema
- ☐ Diffuse vertebral body oedema
- ☒ Posterior element bone marrow oedema
- ☒ Spinous process bone marrow oedema
- ☒ Vertebral corner fat infiltration
- ☒ Syndesmophyte formation
- ☐ All of the above
- ☐ Combination of the pathologies ticked above
- ☐ Other.....